

Differences in Anxiety Levels Between Chewing Gum and Lemon Inhalation Aromatherapy in Hemodialysis Patients

by User id

Submission date: 19-Dec-2023 05:59PM (UTC-0800)

Submission ID: 2262587151

File name: TURNITIN_DYAH_rev.docx (71.85K)

Word count: 3409

Character count: 19337

Differences in Anxiety Levels Between Chewing Gum and Lemon Inhalation Aromatherapy in Hemodialysis Patients

Dyah Restuning Prihati¹, Ulyatusholikah², Maulidta Karunianingtyas Wirawati³

ABSTRACT

End-stage kidney disease requires long-term management and renal replacement therapy such as hemodialysis (HD), peritoneal dialysis, or transplantation. Patients with chronic kidney disease who undergo regular hemodialysis therapy can increase life expectancy; however, hemodialysis therapies can cause stressors leading to psychological problems. This research aims to determine the difference in anxiety levels between chewing gum and lemon inhalation aromatherapy in hemodialysis patients. Furthermore, this research uses a quantitative quasi-experiment research approach with a two-group pre-posttest design, while a purposive sampling technique employs 32 samples at the Hemodialysis Unit of Permata Medika Hospital, Semarang. In addition, data analysis uses the Wilcoxon signed test and the Mann-Whitney test. The research was conducted in July 2023. The research results show that the Wilcoxon test p-value for both interventions was 0.001, and the Mann-Whitney test p-value was 0.494. It can be concluded that there was no difference in anxiety levels between chewing gum and lemon inhalation aromatherapy in hemodialysis patients. The use of non-pharmacological techniques of chewing gum and lemon inhalation aromatherapy can reduce patient anxiety levels during hemodialysis.

Keywords:

Anxiety; Aromatherapy; Hemodialysis; Chewing Gum

1. INTRODUCTION

End-stage kidney disease is a worldwide public health issue because patients must continue to get hemodialysis (HD) for the rest of their lives. In 2016, End-stage kidney disease was ranked fifth on a global scale and is expected to be ranked fifth as the leading cause of death by 2040 [1]. The number of people living with Chronic Kidney Disease (CKD), Acute Kidney Injury (AKI), and Renal Replacement Therapy (RRT) surpasses 850 million, which is more than double the anticipated number of people living with diabetes mellitus and AIDS/HIV worldwide [1]. Over the last two decades, chronic kidney disease has been one of the leading causes of death globally [2]. End-stage renal disease is more likely in patients with a history of diabetes, hypertension, and hypercholesterolemia [3].

The end-stage kidney disease exhibits somatic symptoms that result in reduced health-related quality of life and lifestyle limitations. Ultimately, patients with end-stage kidney disease experience psychological stress [4]. Chronic kidney disease damages the filtering system of the kidneys and nephrons, causing a decrease in the ability to eliminate metabolic waste and a buildup of urea in the blood, which can poison all organs. Long-term care and renal replacement therapy, such as hemodialysis (HD), peritoneal dialysis, or transplantation, are necessary for end-stage kidney disease [5]. Hemodialysis is a procedure for removing blood from the patient's body and circulating it in a machine outside the body (dialyzer) [6]. Kidney hemodialysis therapy takes 4-5 hours for one cycle. Hemodialysis will result in changes in quality of life, so patients need to adapt to their physical condition. Although hemodialysis is the main treatment for end-stage kidney disease, it also causes psychosocial problems. Anxiety is brought on by financial difficulties, therapy-related stress, managing diet, and familial issues in hemodialysis patients [7]. The higher incidence of chronic kidney disease is in line with the increasing costs of hemodialysis therapy the patients must undergo throughout their lives [8].

Patients receiving hemodialysis frequently experience anxiety, which makes it difficult for them to take their medications, manage their fluids, and take care of themselves. Patients with end-stage kidney disease may experience psychological issues such as anxiety. Anxiety manifests physically as headaches, restlessness, perspiration, and palpitations in addition to feelings of stress and tension [7]. Hemodialysis machine alarms, central venous catheter insertion, needle insertion into fistulas, and patient loneliness during hemodialysis are some of the factors that contribute to anxiety in these patients [9]. Accepting one's physical and physiological state as a crucial therapeutic tool for treating pathological, psychological, emotional, and social issues is the first step toward practicing self-care. The treatment of hemodialysis patients cannot be sufficiently effective to produce the intended therapeutic outcomes if patients are not included and self-care activities are not implemented [9]. Chewing sugar-free gum can help hemodialysis patients with dry mouth, but it had no effect on dialysis-related weight gain or thirst. Due to its affordable price and tasty flavor, sugar-free chewing gum has been popular among hemodialysis patients; nonetheless, there is ongoing debate regarding its potential therapeutic benefits [10].

The aromatherapy types commonly used are roses, lavender (*Lavandula*), bergamot, oranges, and lemons. One of the interventions carried out to overcome anxiety is providing aromatherapy (rose, lavender,

lemon). Anxiety activates the sympathetic nervous system and releases stress-related hormones, namely cortisol and adrenaline, which can increase pain levels [11]. For convenience and cost, non-pharmacological treatments like massage, acupuncture, aromatherapy, and relaxation techniques are now being used. Prior studies compared the effects of five drops of either lavender or citrus aurantium essential oils for thirty minutes to the anxiety and agitation of conscious patients admitted to the intensive care unit (ICU) with a placebo control group. Five drops of either lavender or citrus aurantium essential oil were provided to each aromatherapy group, and they were each allowed to inhale for thirty minutes. For 30 minutes, 5 drops of regular saline were administered to the placebo group. Following the intervention, the lavender and citrus aurantium groups experienced much less anxiety than the placebo group. The anxiety levels of the lavender and citrus aurantium groups did not significantly differ from one another. The study's findings demonstrate that aromatherapy with lavender and citrus aurantium helps patients receiving ICU treatment feel less anxious [12]. Patient anxiety is greatly decreased by citrus aurantium. Essential oil of orange aurantium has calming properties, elevates mood, and activates the central nervous system [13]. The mean differences in trait anxiety scores (pre- and post-intervention) do not depend on factors such as gender, age, education level, marital status, reason for starting dialysis, or length of treatment. This is demonstrated by earlier studies that found that the intervention group's four weeks of breathing rose water aromatherapy helped lower anxiety during hemodialysis treatment [7]. Research on two groups, one using an effleurage massage and a combination of essential oils from *Matricaria recutita* L and *Lavandula angustifolia* mill, demonstrated the efficacy of aromatherapy. The control group, meanwhile, was only given effleurage massages. The 14 sessions, which take place once a week, last 30 minutes each. Consequently, there was a noteworthy 30% and 12% decrease in stress and anxiety levels in the intervention group. By contrast, the control group saw a lesser decrease in stress and anxiety levels of 2.6% and 3.3%, respectively [14].

Preliminary studies in the Hemodialysis Room at Permata Medika Hospital Semarang in March 2023 on 10 intra-hemodialysis patients show that two patients experienced mild anxiety with a score of 30-40 (State-Trait Anxiety Inventory questionnaire). They underwent hemodialysis for the first time, accompanied by family. Furthermore, three patients experienced moderate anxiety with a score of 41-50 and no family to accompany them. Four patients experienced severe anxiety with a score of 51-80 due to fear of hemodialysis complications they had experienced. Finally, one patient was not anxious since a family member was accompanying, and this was not the first hemodialysis procedure. This condition can affect the psychological condition of chronic kidney disease patients when undergoing hemodialysis therapy. In addition, the nurses need to provide an intervention to reduce anxiety and provide a sense of comfort for patients undergoing hemodialysis to make chronic kidney disease patients carry out the therapy more easily.

2. RESEARCH METHOD

The research method used in this study is a quasi-experimental two-group pre-posttest design. The research respondents are 47 hemodialysis patients at Permata Medika Hospital, Semarang, divided into 2 intervention groups: chewing gum and lemon inhalation aromatherapy. The inclusion criteria for respondents include the general condition of the respondent being stable, having no problems with the mouth or smell, having no allergies, and having anxiety. This study uses the State-Trait Anxiety Inventory (STAI) anxiety level questionnaire by Spielberger (1983) [15], modified in Indonesian with correlation index results ranging from 0.196 to 0.664 and a reliability of 0.890. Research data were collected in July 2023. The research was carried out in the hemodialysis unit at Permata Medika Hospital, Semarang. Researchers conducted a pre-test using the STAI questionnaire on respondents. The intervention is given to each group after 30 minutes of hemodialysis. The patients in the intervention group chew 2 pieces of xylitol gum (not to be swallowed) for ± 10 minutes. Meanwhile, the other group of patients get aromatherapy for 30 minutes every time they undergo hemodialysis. Lemon aromatherapy is presented as tissue or gauze that has been dripped with oil (3 drops or 0.3 ml) and placed right next to the respondent's pillow at a distance of 20-30 cm from the respondent's nose. Respondents are asked to inhale the aroma during the hemodialysis process. The intervention is given every time the patients undergo hemodialysis at 4 meetings for two weeks. After two weeks, a post-test is conducted to measure the patient's anxiety level with the STAI questionnaire after 30 minutes of intervention.

Univariate data analysis includes respondent characteristics (age, gender, education). These characteristics are displayed in a frequency distribution. Bivariate data analysis in the study uses the Wilcoxon test to determine hemodialysis patients' anxiety before being given intervention in each gum-chewing group and lemon inhalation aromatherapy group. The Mann-Whitney test is employed to determine the difference in anxiety levels between chewing gum and lemon inhalation aromatherapy in hemodialysis patients. This research has received approval from the ethics committee of Widya Husada University Semarang No. 68/EC-LPPM/UWHS/VIII-2023.

3. RESULTS AND DISCUSSION

3.1. Overview of respondents

Table 1 shows the frequency distribution of respondents based on the characteristics of hemodialysis respondents. Most patients are 45-59 years old, namely 19 (59.4%) people. There is a minimum and maximum age of 32-70 years and a mean value of 51.66 years with the largest gender being female, totaling 23 (71.9%) people. In addition, most patients had elementary school education, namely 11 (34.4%) people.

Table 1. Frequency Distribution of Respondents Based on Characteristics of Hemodialysis Respondents

No.	Characteristic	Frequency	Percentage (%)
1.	Age		
	30-44	6	18.75
	45-59	19	59.4
	60-69	6	18.75
	70	1	3.1
2	Sex		
	Male	9	28.1
	Female	23	71.9
3	Education		
	No Educational Background	2	6.25
	Primary School	11	34.4
	Junior High School	7	21.9
	Senior High School	10	31.25
	D3	1	3.1
	S1	1	3.1
	Total	32	100

An individual's maturity in dealing with anxiety is very important. This is influenced by a person's coping mechanisms; the more mature an individual is, the better their readiness to deal with anxiety. In addition, the older the person, the greater the risk that a person will face regarding bodily health, especially the kidneys. Hemodialysis treatment over a long period of time can cause complications and anxiety in patients with chronic kidney disease. However, as a person gets older, it will influence their thinking patterns in a positive direction. Therefore, if a person is faced with an anxious condition, it can be easily resolved. Chronic kidney disease is more common in women, but the prevalence of end-stage kidney disease who receive kidney replacement therapy is predominantly in men. Causative factors include employment, low adherence to treatment, and low utilization of health facilities [16]. Men have unhealthy lifestyles such as irregular eating patterns, consuming minimal alcohol, smoking, and having higher creatinine levels compared to women.

A person's emotional maturity and capacity to handle a variety of situations improve with age, although anxiety can strike at any stage of life. Any number of coping strategies may be the source of this for any given person. Males tend to be more energetic and exploratory, while women are more sensitive, according to previous studies, and women are more worried than males. Aside from that, women place a great deal of importance on appearance. However, physical changes such as changes in skin tone, dryness, itching, and hair loss that come with chronic renal failure alter a person's perception of their own attractiveness. Hemodialysis patients in Egypt experience anxiety and depression, and a number of sociodemographic risk factors include smoking, a sedentary lifestyle, female gender, and higher hemodialysis duration [17]. This contradicts earlier studies that found men to have higher levels of anxiety and despair than women [18].

Anxiety is a common occurrence because a lower level of knowledge is associated with a decreased capacity to identify and explain novel difficulties. Reasoning and absorbing new information—including describing novel problems—will be made easier with a higher degree of education. One can enhance their quality of life by obtaining a high degree of education. A person's positive behavior is influenced by their educational attainment. This is because, particularly for those receiving hemodialysis, education may be one of the elements contributing to an improved quality of life. When a patient is unsure of the surgery they will have and worries about the outcome of their therapy, anxiety levels may be higher during the first few hours of the process. Patients learn the process and start to adjust after receiving hemodialysis. The patient is already in the acceptance stage, thus the longer they receive hemodialysis therapy, the lower their anxiety level will be. Patients on dialysis are affected by anxiety 1.5 times more often [19].

3.2. The Effect of Chewing Gum on Anxiety in Hemodialysis Patients

Table 2 shows negative rank values for 15 respondents and tie values for 1 respondent, meaning there was a decrease in anxiety scores between the pretest and posttest, and 1 respondent had the same score

between the pretest and posttest. The p-value is 0.001 ($p < 0.05$), meaning that H_a is accepted. There was an influence on the anxiety level of hemodialysis patients before and after being given the chewing gum intervention.

Table 2. Effect of Chewing Gum on Anxiety in Hemodialysis Patients

Posttest – Pretest	N	Mean Rank	P-Value
Negative Ranks	15	8.00	
Positive Ranks	0	.00	0.001
Ties	1		
Total	16		

The safest gum to chew is sugar-free gum that contains xylitol. Xylitol is a natural sweetener that helps reduce stress and anxiety [20]. It also helps reduce cravings for sweet foods, plaque, and cavities [21]. However, it should be noted that chewing gum excessively every day can cause jaw pain and headaches. Chewing gum can reduce anxiety levels in certain situations due to the release of endorphins that are triggered when chewing gum. Candy containing xylitol is a natural sweetener that can help relax the body. Meanwhile, chewing gum containing chamomile is an herbal medicine that contains sedatives. These two ingredients combined can help reduce feelings of stress and anxiety. Chewing gum helps calm and focus the mind, reduces symptoms of depression, increases self-confidence by reducing anxiety, and provides a feeling of calm and relaxation. Additionally, chewing gum increases saliva production, which can help reduce tooth decay and freshen breath [22]. Chewing gum can reduce salivary cortisol levels. In this case, the chewing movement can reduce anxiety because this movement can reduce muscle tension and energy produced through chewing. This process occurs at the beginning of digestion and increases saliva flow, which causes cortisol levels to decrease and is followed by a decrease in anxiety levels.

One hundred students who were given the intervention of chewing gum prior to the exam in order to reduce exam stress and improve exam performance were the subjects of the study. Gum chewing helps ease anxiety and tension. Gum chewing helps to enhance short-term memory. However, chewing gum can lead to jaw stress, headaches, and digestive issues, therefore it is not advised for pupils who struggle with concentration over the long run. Chewing gum to relieve anxiety has the potential to develop into a habit that could become addictive [23]. Students who chewed gum for 20 minutes a day reported less stress overall, with the biggest benefits occurring when they had a heavy workload [24]. While chewing gum reduces beforehand anxiety without raising the risk of pulmonary aspiration, it does aid women experiencing gynecological preoperative anxiety and dry mouth during the preoperative fasting period [25].

3.3. The Effect of Lemon Inhalation Aromatherapy on Anxiety in Hemodialysis Patients

Table 3 shows negative rank values for 14 respondents and tie values for 2 respondents, meaning there was a decrease in anxiety scores between the pretest and posttest, and 2 respondents had the same scores between the pretest and posttest. The p-value is 0.001 ($p < 0.05$), meaning that H_a is accepted. There was an influence on the anxiety level of hemodialysis patients before and after being given the lemon aromatherapy inhalation intervention.

Table 3. Effect of Lemon Inhalation Aromatherapy on Anxiety in Hemodialysis Patients

Posttest – Pretest	N	Mean Rank	P-Value
Negative Ranks	14	7.50	
Positive Ranks	0	0.00	0.001
Ties	2		
Total	16		

Aromatherapy is a simple and effective method for dealing with anxiety. Aromatherapy involves the use of the psychological, physiological, and medicinal effects of essential oils to improve the patient's condition. Aromatherapy can regulate emotions positively, control anxiety, reduce depression, and stimulate or strengthen memory. The particles in aromatherapy stimulate the olfactory nerves and then begin to be sent to the limbic system. Stimulation sent to the limbic system can stimulate the hypothalamus and then stimulate the vagus nerve. The vagus nerve conveys its parasympathetic signals to the heart, which then reduces the frequency and contractility of the heart. Hence, aromatherapy data provides a feeling of relaxation and calm in patients with anxiety [26].

The study results show that the patient's anxiety can be overcome with lemon inhalation aromatherapy intervention. There were tie scores of 2 respondents who did not have a change in category in the pretest and posttest in providing this intervention even though there was a decrease in scoring values. This was due to the length of time hemodialysis was carried out routinely and continuously, making the patient bored and anxious. Meanwhile, there was a negative rank value of 14 respondents and a positive rank of 0, meaning that there was a decrease in the anxiety level value from pretest to posttest in a positive direction. One of the substances contained in lemon is linalool which is useful for stabilizing the nervous

system to have a calming effect and reduce a person's anxiety level. Factors that influence reducing anxiety when giving aromatherapy include the content contained in lemon, the accuracy of the method of administration, the duration of administration, the quality of the essential oil, and the right dose so as not to cause side effects such as nausea and dizziness.

Reducing preoperative anxiety in individuals who are waiting for surgery can be achieved with the use of aromatherapy. When it came to reducing preoperative anxiety, shorter sessions (≤ 20 minutes) of aromatherapy performed better than longer sessions (> 20 minutes) [27]. Physiological measurements of vital indicators, such as blood pressure, heart rate, respiration rate, pulse rate, and saturation, can be used to objectively determine the degree of anxiety following lavender oil inhalation [28]. Acute coronary syndrome patients experience anxiety, an unpleasant emotion that raises the heart's oxygen demand. Elevations in anxiety levels have been shown to alter blood pressure, pulse rate, respiratory rate, and myocardial oxygen consumption via raising plasma concentrations of adrenaline and norepinephrine [26].

3.4. Differences in Anxiety Levels Between Chewing Gum and Lemon Inhalation Aromatherapy in Hemodialysis Patients

Table 3 shows that the Mann-Whitney test obtained a p-value of 0.494 ($p > 0.05$), which means that H_a is rejected, and H_o is accepted. In addition, there was no difference in anxiety levels between chewing gum and inhaling lemon aromatherapy in hemodialysis patients.

Table 4. Differences in Anxiety Levels Between Chewing Gum and Lemon Inhalation Aromatherapy in Hemodialysis Patients

Anxiety Level	N	P-Value
Chewing Gum	16	0.494
Inhaling Lemon Aromatherapy	16	
Total	32	

Moreover, chewing gum and lemon aromatherapy inhalation interventions are effective for hemodialysis patients who experience anxiety. This is because the two interventions are relatively easy to implement and have cheap ingredients and can be found anywhere. However, comparing both interventions shows neither of which is superior.

4. CONCLUSION

The study results show no difference in the level of anxiety between chewing gum and lemon inhalation aromatherapy for hemodialysis patients. The Mann-Whitney test shows the p-value is 0.494 ($P > 0.05$). The limitations of this study show that researchers were unable to explore in more depth other factors that could cause chronic kidney disease patients to experience anxiety when undergoing hemodialysis therapy. The results of this research can be used as evidence-based nursing practice for nurses in hemodialysis rooms regarding the use of non-pharmacological techniques by chewing gum and lemon inhalation aromatherapy on patient anxiety levels.

Differences in Anxiety Levels Between Chewing Gum and Lemon Inhalation Aromatherapy in Hemodialysis Patients

ORIGINALITY REPORT

10%

SIMILARITY INDEX

6%

INTERNET SOURCES

5%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Mohsen Soleimani, Leyla Sadat Kashfi, Majid mirmohamadkhani, Ali Asghar Ghods. "The effect of aromatherapy with peppermint essential oil on anxiety of cardiac patients in emergency department: A placebo-controlled study", *Complementary Therapies in Clinical Practice*, 2022
Publication 1%

- 2** Zahra Karimzadeh, Mansooreh Azizzadeh Forouzi, Elham Rahiminezhad, Mehdi Ahmadinejad, Mahlagha Dehghan. "The Effects of Lavender and Citrus aurantium on Anxiety and Agitation of the Conscious Patients in Intensive Care Units: A Parallel Randomized Placebo-Controlled Trial", *BioMed Research International*, 2021
Publication 1%

- 3** www.scilit.net
Internet Source 1%

- 4** link.springer.com
Internet Source

1 %

5

marketdesigner.blogspot.com

Internet Source

1 %

6

repository.unsoed.ac.id

Internet Source

1 %

7

www.zora.uzh.ch

Internet Source

1 %

8

Guillermo García García, Arpana Iyengar, François Kaze, Ciara Kierans, Cesar Padilla-Altamira, Valerie A. Luyckx. "Sex and gender differences in chronic kidney disease and access to care around the globe", *Seminars in Nephrology*, 2022

Publication

<1 %

9

Submitted to Clarkston Community Schools

Student Paper

<1 %

10

Submitted to St George's Hospital Medical School

Student Paper

<1 %

11

Submitted to Queen's University of Belfast

Student Paper

<1 %

12

health-plaza.info

Internet Source

<1 %

13

Submitted to Laureate Higher Education Group

Student Paper

<1 %

14

Fulya Türk, Ayşe Nur Katmer. "A study on the effectiveness of coping with test anxiety program based on cognitive-behavioral approach", International Journal of Evaluation and Research in Education (IJERE), 2019

Publication

<1 %

15

Sezer Avcı, Tulay Ortabag, Hasan Ulusal, Seyithan Taysi. "The Effect of Domestic Violence During Pregnancy on Cortisol Hormone Release, Breastfeeding, and Newborn", Journal of Interpersonal Violence, 2022

Publication

<1 %

16

[archive.org](https://www.archive.org)

Internet Source

<1 %

17

[business.dailytimesleader.com](https://www.business.dailytimesleader.com)

Internet Source

<1 %

18

John M. Harris. "Can internet-based continuing medical education improve physicians' skin cancer knowledge and skills?", Journal of General Internal Medicine, 01/2001

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off